

11PP

RECEIVED  
OPPT NCIC


OPPT-2002-0029-0027

2002 JUL 29 PM 3:19




## Advantages of a Two-Generation Mysid Protocol


Charles L. McKenney, Jr.  
Office of Research and Development  
National Health and Environmental  
Effects Research Laboratory  
Gulf Ecology Division  
Gulf Breeze, FL 32561



### *Americamysis bahia*




### *Americamysis bahia* Life-Cycle




## Summary of Mysid Life-Cycle Exposures


### ↓ YOUNG PRODUCTION





### Evaluating effects of potential endocrine-disrupting compounds - a life-cycle context




# Effect of Juvenile Hormone Agonists (JHA)






ORGANISM	EFFECT CONCENTRATION METHOPRENE (mg/l)
RAT	210
FISH	
RAINBOW TROUT	106
CHANNEL CATFISH	100
BLUEGILL	4.62
COHO SALMON	86
MUMMICHOOG	125
MOSQUITOES	
<i>Culex pipiens</i>	0.1-1.0
<i>C. tritaeniorhynchus</i>	0.1
<i>Aedes nigromaculatus</i>	0.1
<i>A. quadrimaculatus</i>	0.01
<i>A. taeniorhynchus</i>	0.0001
CRUSTACEANS	
<i>Palaemonetes pugio</i>	0.008
<i>Rhithropanopeus harrisa</i>	0.1
<i>Daphnia magna</i>	0.1
<i>Mysidopsis bahia</i>	0.002




# Endocrine Regulation of Crustacean Reproduction and Growth



From Charmantier et al., 1997


# *Americanysis bahia* vs pyriproxyfen exposures



[Ecdysteroids] of day 1, brood 3  
females vs pyriproxyfen



# TRANSGENERATIONAL EFFECTS ON MYSIDS

## 1st. GENERATION EXPOSED





Altered development, growth and reproduction of  
Crustaceans by EDGs would restrict their role as  
dominant secondary producers

